






A new four-pored *Amphisbaena* Linnaeus, 1758 (*Amphisbaenia*, *Amphisbaenidae*) from the north of Espinhaço Mountain Range, Brazil

Síria Ribeiro^{1,2}, Alfredo P. Santos Jr^{1,2}, Isabelly G. Martins^{1,2}, Elaine C. S. Oliveira³, Roberta Graboski⁴, Thiago Barbosa Da Silveira⁵, Matheus H. M. Benício⁶, Wilian Vaz-Silva⁷

¹ Universidade Federal do Oeste do Pará, Programa de Pós-Graduação em Biodiversidade, 68040-255, Santarém, Pará, Brazil

² Laboratório de Ecologia e Comportamento Animal, Universidade Federal do Oeste do Pará, 68040-255, Santarém, Pará, Brazil

³ Instituto Chico Mendes de Conservação da Biodiversidade, Coordenação Regional Oeste do Pará, 68040-000, Santarém, Pará, Brazil

⁴ School of Zoology, Faculty of Life Sciences, the Steinhardt Museum of Natural History, Tel-Aviv University, Tel-Aviv, Israel

⁵ Tecsan – Tecnologia e Saneamento Ltda – Rua Dr. José Peroba, 275, Ed. Metrópolis Empresarial, Sala 201/202, Stiep, 41770-235, Salvador, Bahia, Brazil

⁶ Bahia Mineração S/A, Rodovia BA 156, s/n, Projeto Pedra de Ferro, 46400-000, Caetité, Bahia, Brazil

⁷ Pontifícia Universidade Católica de Goiás, Escola de Ciências Médicas e da Vida, Centro de Estudos e Pesquisas Biológicas, and Programa de Pós-Graduação em Ciências Ambientais e Saúde, 74605-010, Goiânia, Goiás, Brazil

Corresponding author: Síria Ribeiro (siherp@hotmail.com)



Academic editor: Uri García-Vázquez

Received: 4 March 2024

Accepted: 18 July 2024

Published: 24 September 2024

ZooBank: <https://zoobank.org/77F68F11-081E-4693-94BD-F211FFCFED62>

Citation: Ribeiro S, Santos Jr AP, Martins IG, Oliveira ECS, Graboski R, Barbosa Da Silveira T, Benício MHM, Vaz-Silva W (2024) A new four-pored *Amphisbaena* Linnaeus, 1758 (*Amphisbaenia*, *Amphisbaenidae*) from the north of Espinhaço Mountain Range, Brazil. ZooKeys 1213: 1–27. <https://doi.org/10.3897/zookeys.1213.122265>

Copyright: © Síria Ribeiro et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

A new species of *Amphisbaena* is described from the north of Espinhaço Mountain Range, municipality of Caetité, state of Bahia, Brazil. *Amphisbaena amethysta* **sp. nov.** can be distinguished from its congeners by the following combination of characters: (1) snout convex in profile, slightly compressed not keeled; (2) pectoral scales arranged in regular annuli; (3) four precloacal pores; (4) distinct cephalic shields; (5) 185–199 dorsal half-annuli; (6) 13–16 caudal annuli; (7) conspicuous autotomic site between 4th–6th caudal annuli; (8) 16–21 dorsal and ventral segments at midbody; (9) 3/3 supralabials; (10) 3/3 infralabials; and (11) smooth and rounded tail tip. The new species is the 71st species of genus with four precloacal pores, and the 22nd species from the Caatinga morphoclimatic domain. The identification of *Amphisbaena amethysta* **sp. nov.** indicates that the reptile fossorial fauna in the Espinhaço Mountain Range region is far from being completely known and that it may harbour a much greater diversity of endemic taxa.

Key words: Morphology, new species, phylogeny, taxonomy

Introduction

In the Caatinga 21 species of *Amphisbaena* have been recorded to date, with most being fully restricted to this morphoclimatic domain (Ribeiro and Eleutério 2023). Of these, four have wide distributions in Caatinga: *A. alba* Linnaeus, 1758, *A. lumbricalis* Vanzolini, 1996, *A. pretrei* Duméril & Bibron, 1839, and *A. vermicularis* Wagler, 1824, while nine have been found in a small number of localities: *A. anomala* (Barbour, 1914), *A. arenaria* Vanzolini, 1991, *A. bahiana* Vanzolini, 1964, *A. carvalhoi* Gans, 1965, *A. frontalis* Vanzolini, 1991, *A. fuliginosa*

Linnaeus, 1758, *A. hastata* Vanzolini, 1991, *A. heathi* Schmidt, 1936, and *A. ignatiana* Vanzolini, 1991. In addition, five species are known only from the type locality: *A. arda* Rodrigues, 2003, *A. caetitensis* Almeida, Freitas, Silva, Valverde, Rodrigues, Pires & Mott, 2018, *A. longinqua* Teixeira Junior, Dal Vechio, Recoder, Cassimiro, Sena & Rodrigues, 2019, *A. mongoyo* Teixeira Junior, Dal Vechio, Recoder, Cassimiro, Sena & Rodrigues, 2019, and *A. uroxena* Mott, Rodrigues, Freitas & Silva, 2008 (Vanzolini 1992; Teixeira Jr et al. 2014; Costa et al. 2015; Almeida et al. 2018; Ribeiro and Eleutério 2023).

In the last two decades, six species of *Amphisbaena* have been identified in the high-altitude areas of the Espinhaço Mountain Range in Bahia and Minas Gerais: *A. bahiana*, *A. longinqua*, *A. metallurga* Costa, Resende, Teixeira Jr, Dal Vechio & Clemente, 2015, *A. mongoyo*, *A. uroxena*, and *A. caetitensis* (Costa et al. 2015; Almeida et al. 2018; Teixeira Junior et al. 2019). Four of which are known to be closely related phylogenetically (Teixeira Junior et al. 2019), while the position of the *A. metallurga* awaits analysis; and that of *A. caetitensis* awaits resolution of the phylogenetic relationship (Almeida et al. 2018). Such clustering points to a probable centre of endemism for this type of geological formation.

During a faunal rescue in the south of state of Bahia, Brazil, conducted as part of environmental activities carried by BAMIN (Bahia Mineração mining company), specimens of a species of *Amphisbaena* with four precloacal pores were collected, which could not be identified as belonging to any known amphisbaenid species. Accordingly, it was concluded that they represented a new taxon, which is described below.

Materials and methods

We analysed 48 specimens of non-identified *Amphisbaena* from the municipality of Caetité, state of Bahia, Brazil. The type series was deposited in the herpetological collection from Pontifícia Universidade Católica de Goiás (CEPB), municipality of Goiânia, state of Goiás, Brazil. For morphological comparisons we used data from 370 analysed specimens of *Amphisbaena* (see list S11). Additional morphological data was taken from the literature. The taxonomy follows the nomenclature of Guedes et al. (2023) for species (but not including subspecies). Nomenclature for the cephalic scales and for meristic data follow Gans and Alexander (1962). Morphometric data follow the methodology of Perez et al. (2012). Head scale morphometric data were taken with digital callipers (precision 0.01 mm) on the right side of specimens. Head length was measured from the tip of the rostral shield to the anterior margin of the first dorsal half-annulus; and ventral length of head was measured from tip of rostral shield to the anterior margin of first body annulus. Snout-vent length was measured with the aid of a nylon line and subsequently measured with a millimetre rule. Counts for dorsal and ventral half-annuli were made on the right side of each specimen. Incomplete half-annuli were not included in the total count. Observations on specimen and annuli counts were made with the aid of a stereomicroscope. Infralabials, supralabials, and parietal shields variations are treated as “right/left”.

Our molecular data matrix comprises 81 terminals for six genes: three mitochondrial genes (16S – 16S Large Subunit Ribosomal RNA gene; 12S – Small Subunit Ribosomal RNA gene; and *nd2* – NADH Dehydrogenase 2 gene) and

three nuclear genes (*c-mos* – Oocyte maturation factor Mos; *bdnf* – Brain-derived neurotrophic factor; and *rag1* – Recombinant activating gene 1). We sequenced four new DNA fragments (two for 12S and two for 16S) for two specimens of *Amphisbaena amethysta* sp. nov. (see Suppl. material 1: table S1). We also included sequences available in GenBank (<https://www.ncbi.nlm.nih.gov>) for 79 species of *Amphisbaena*. We rooted our phylogenetic tree using Lacertidae (*Lacerta media* Lantz & Cyrén, 1920).

DNA was extracted from liver tissue using the PureLink extraction kit (Invitrogen, Massachusetts, USA), following the manufacturer's protocol. Sequences were amplified by Polymerase Chain Reaction (PCR) using the primers 12S and 16S as described by Graboski et al. (2023) following the amplification protocols described in Kearney and Stuart (2004) and Mott and Vieites (2009). Amplified fragments were purified with shrimp alkaline phosphatase and exonuclease I (GE Healthcare, Piscataway, NJ, USA). Both strands were sequenced on an Applied Biosystems 3500 Series Genetic Analyzer (Thermo Fisher Scientific, USA) at Laboratório de Genética e Biodiversidade da Universidade Federal de Goiás, Brazil. Both strands were quality-checked and, when necessary, edited manually. Consensus for both strands was generated using Geneious Prime 2022.1.1 (<https://www.geneious.com>).

Sequences were aligned using MAFFT 1.3.6 (Katoh and Standley 2013) through a plugin implemented in Geneious Prime. The 16S and 12S sequences were aligned under the E-INS-I algorithm, while *nd2* and nuclear genes were aligned under the G-INS-i algorithm. We used default parameters for gap opening and extension. The protein-coding gene alignments were visually checked using Geneious Prime to verify that all sequences follow the correct reading frame. All genes were concatenated using Geneious Prime.

We used PartitionFinder 2 (Lanfear et al. 2016) to identify the combined best-fitting of partitioning schemes and models of molecular evolution. Our input matrix was divided in 14 partitions (coding genes were partitioned by codon positions and each rRNA was analysed as a separate partition) and was analysed using the greedy option. We performed a run allowing the program to select (using the Akaike Information Criterion with correction: AIC) for molecular evolution models implemented on RAxML (models GTR and GTR+G). We performed a maximum likelihood (ML) analysis using RAxML 8.2.3 (Stamatakis 2014). The ML tree was estimated using the RAxML algorithm that conducts a rapid bootstrap analysis and searches for best scoring ML tree in the same run (option -f a). We ran 1000 bootstrap replicates, and the best scoring ML tree was estimated 200 times using as a starting tree each fifth bootstrap tree. We also calculate uncorrected genetic distance (p-distance) using PAUP 4.0 (Swofford 2003). We considered only bootstrap values above 70% as a strong support.

Results

Amphisbaena amethysta sp. nov.

<https://zoobank.org/E34029C4-6E1A-4E87-89C7-99E70AC1B95A>

Figs 1–4

Type material. Holotype: • male; CEPB 2311; municipality of Caetité, state of Bahia, BRAZIL; [14°21'31"S, 42°32'19"W; 1012 m above sea level (a.s.l.)]; collected on 1 November 2022 by Faunal Rescue Team Tecsam (F. Santos, P. Belufi, and

G. Nascimento). **Paratypes:** • All from Caetité, Bahia, Brazil; collected by Faunal Rescue Team Tecsam (R. Assunção, A. Hirota, T. Silveira, F. Santos, P. Belufi, and G. Nascimento) • Female; CEPB 2301; (14°21'53"S, 42°32'20"W; 1013 m a.s.l.); 8 June 2022 • Male; CEPB 2302; (14°01'16"S, 42°31'06"W; 1082 m a.s.l.); 13 October 2021 • Female; CEPB 2303; (14°21'50"S, 42°32'20"W; 1013 m a.s.l.); 26 May 2022 • Female; CEPB 2308; (14°19'49"S, 42°32'44"W; 1011 m a.s.l.); 8 September 2022 • Female; CEPB 2327; (14°19'52"S, 42°32'42"W; 1011 m a.s.l.); 1 October 2022 • Female; CEPB 2331; (14°19'45"S, 42°32'45"W; 1011 m a.s.l.); 30 August 2022 • Male; CEPB 2346; (14°22'08"S, 42°32'17"W; 923 m a.s.l.); 6 October 2022 • Male; CEPB 2379; (14°21'27"S, 42°32'20"W; 1033 m a.s.l.); 26 January 2023 • Female; CEPB 2381; (14°21'27"S, 42°32'20"W; 1033 m a.s.l.); 26 January 2023.

Referred specimens. All from Caetité, Bahia, BRAZIL; collected by Faunal Rescue Team Tecsam (R. Assunção, A. Hirota, T. Silveira, F. Santos, P. Belufi, and G. Nascimento) • Female; CEPB 2298; (14°21'53"S, 42°32'20"W; 1013 m a.s.l.); 8 June 2022 • Female; CEPB 2299; (14°21'53"S, 42°32'20"W; 1013 m a.s.l.); 8 June 2022 • Male; CEPB 2300; (14°21'51"S, 42°32'21"W; 1011 m a.s.l.); 25 May 2022 • Female; CEPB 2304; (14°21'53"S, 42°32'20"W; 1013 m a.s.l.); 8 June 2022 • Female; CEPB 2305; (14°21'53"S, 42°32'20"W; 1013 m a.s.l.); 8 June 2022 • Female; CEPB 2306; (14°21'53"S, 42°32'20"W; 1013 m a.s.l.); 13 August 2022 • Male; CEPB 2307; (14°21'23"S, 43°32'07"W; 972 m a.s.l.); 23 November 2022 • Undetermined sex; CEPB 2309; (14°19'46"S, 42°43'20"W; 1011 m a.s.l.); 1 September 2022 • Female; CEPB 2310; (14°19'55"S, 42°32'44"W; 1011 m a.s.l.); 8 October 2022 • Male; CEPB 2312; (14°19'56"S, 42°32'43"W; 1011 m a.s.l.); 12 October 2022 • Female; CEPB 2313; (14°22'08"S, 42°32'17"W; 9215 m a.s.l.); 6 October 2022 • Female; CEPB 2314; (14°22'08"S, 42°32'17"W; 925 m a.s.l.); 6 October 2022 • Female; CEPB 2315; (14°22'08"S, 42°32'17"W; 925 m a.s.l.); 6 October 2022 • Female; CEPB 2316; (14°19'47"S, 42°32'43"W; 1011 m a.s.l.); 1 September 2022 • Female; CEPB 2317; (14°19'55"S, 43°32'44"W; 1011 m a.s.l.); 1 October 2022 • Female; CEPB 2318; (14°19'46"S, 42°32'43"W; 1011 m a.s.l.); 1 September 2022 • Female; CEPB 2319; (14°20'37"S, 42°32'13"W; 1076 m a.s.l.); 3 August 2022 • Female; CEPB 2320; (14°20'38"S, 42°32'12"W; 1013 m a.s.l.); 2 June 2022 • Female; CEPB 2321; (14°19'53"S, 42°32'43"W; 1011 m a.s.l.); 29 September 2022 • Female; CEPB 2322; (14°14'28"S, 42°32'47"W; 842 m a.s.l.); 17 September 2022 • Female; CEPB 2323; (14°19'49"S, 42°32'17"W; 1062 m a.s.l.); 13 August 2022 • Male; CEPB 2324; (14°19'48"S, 42°32'43"W; 1011 m a.s.l.); 6 September 2022 • Female; CEPB 2325; (14°19'55"S, 42°32'44"W; 1011 m a.s.l.); 1 October 2022 • Male; CEPB 2326; (14°20'52"S, 43°32'16"W; 1053 m a.s.l.); 13 August 2022 • Female; CEPB 2328; (14°21'53"S, 42°32'22"W; 1011 m a.s.l.); 28 September 2022 • Male; CEPB 2329; (14°19'47"S, 42°32'43"W; 1011 m a.s.l.); 1 September 2022 • Male; CEPB 2330; (14°19'55"S, 43°32'44"W; 1011 m a.s.l.); 1 October 2022 • Male; CEPB 2332; (14°20'50"S, 42°32'16"W; 1057 m a.s.l.); 13 August 2022 • Female; CEPB 2333; (14°14'28"S, 42°32'47"W; 842 m a.s.l.); 17 September 2022 • Female; CEPB 2334; (14°19'46"S, 42°32'43"W; 1011 m a.s.l.); 1 September 2022 • Female; CEPB 2336; (14°15'11"S, 45°32'16"W; 987 m a.s.l.); 3 August 2022 • Female; CEPB 2337; (14°20'38"S, 45°32'12"W; 1076 m a.s.l.); 3 August 2022 • Female; CEPB 2338; (14°21'31"S, 45°32'16"W; 1010 m a.s.l.); 13 August 2022 • Female; CEPB 2339; (14°14'28"S, 42°32'47"W; 842 m a.s.l.); 17 September 2022 • Male; CEPB 2356; (14°21'27"S, 42°32'21"W; 1033 m a.s.l.); 1 November 2022 • Male; CEPB 2379; (14°21'27"S, 42°32'20"W; 1033 m a.s.l.); 26 January 2023 • Female; CEPB 2380;




Figure 1. *Amphisbaena amethysta* sp. nov. (Holotype, CEPB 2311) **A** dorsal **B** lateral, and **C** ventral views of head. Scale bar: 3 mm.



Figure 2. Tail of *Amphisbaena amethysta* sp. nov. (Holotype, CEPB 2311). Detail of the autotomic site and four precloacal pores. Scale bar: 3 mm.

(14°21'27"S, 42°32'20"W; 1033 m a.s.l.); 26 January 2023 • Female; CEPB 2381; (14°21'27"S, 42°32'20"W; 1033 m a.s.l.); 26 January 2023.

Diagnosis and comparisons with other south American amphisbaenians.

Amphisbaena amethysta sp. nov. is a medium-sized amphisbaenid (258 mm maximum snout-vent length), and can be distinguished from its congeners by the following combination of characters (see details in Table 1): (1) snout convex in profile view, slightly compressed not keeled; (2) pectoral scales arranged in regular annuli; (3) four precloacal pores; (4) distinct cephalic shields; (5) 185–199 dorsal half-annuli; (6) 13–16 caudal annuli; (7) conspicuous autotomic site between 4th–6th caudal annuli; (8) 16–21 dorsal and ventral segments at midbody; (9) 3/3 supralabials; (10) 3/3 infralabials; and (11) smooth and rounded tail tip.

Amphisbaena amethysta sp. nov. differs from *Amphisbaena acrobeles* (Ribeiro, Castro-Mello & Nogueira, 2009), *A. bilabialata* (Stimson, 1972), *A. kingi* (Bell, 1833), *A. anomala*, *Mesobaena huebneri* Mertens, 1925; *M. rhachicephala* Hoogmoed, Pinto, Rocha & Pereira, 2009; and all *Leposternon* species, mainly in having the snout convex in profile view, slightly compressed not keeled (vs snout hardly compressed forming a sharp and prominent keel or snout depressed shovel-like). Differs still from *A. anomala* and all *Leposternon* species by having pectoral scales arranged in regular annuli (vs pectoral scales with an irregular form, and dermal annuli not regularly arranged).

Amphisbaena amethysta sp. nov. presents four precloacal pores, differing from all *Amphisbaena* species except *A. acangaoba* Ribeiro, Gomides & Costa, 2020, *A. alba*, *A. albocingulata* Boettger, 1885, *A. angustifrons* Cope, 1861, *A. arda*, *A. arenaria*, *A. arenicola* Perez & Borges-Martins, 2019, *A. bahiana*, *A. bakeri* Stejneger, 1904, *A. barbouri* Gans & Alexander, 1962, *A. bedai* (Vanzolini, 1991), *A. bolivica* Mertens, 1929, *A. borellii* Peracca, 1897, *A. brasiliana* (Gray, 1865), *A. caeca* Cuvier, 1829, *A. camura* Cope, 1862, *A. carlgansi* Thomas & Hedges, 1998, *A. carioca* Rocha, Barros-Filho & Sluys, 2023, *A. carvalhoi*, *A. caudalis* Cochran, 1928, *A. cayemite* Thomas & Hedges, 2006, *A. cegei* Montero, Sáfadez, Álvarez, 1997, *A. cubana* Gundlach & Peters, 1879, *A. cuiabana* (Strüssmann & Carvalho, 2001), *A. cunhai* Hoogmoed & Ávila-Pires, 1991,




Figure 3. *Amphisbaena amethysta* sp. nov. in life (not identified specimen of type series) **A** lateral view and **B** dorsal. Photograph by T.B.S.

A. darwini, *A. elbakyanae* Torres-Ramírez, Angarita-Sierra & Vargas-Ramírez, 2021, *A. fenestrata* (Cope, 1861), *A. frontalis*, *A. gonavensis* Gans & Alexander, 1962, *A. gracilis* Strauch, 1881, *A. hastata*, *A. heathi*, *A. hogei* Vanzolini, 1950, *A. hoogmoedi* Oliveira, Vaz-Silva, Santos-Jr, Graboski, Teixeira Jr, Dal Vechio & Ribeiro, 2018, *A. hyporissor* Thomas, 1965, *A. innocens* Weinland, 1862, *A. kingi*, *A. kraoh* (Vanzolini, 1971), *A. leali* Thomas & Hedges, 2006, *A. lumbricalis*, *A. manni* Barbour, 1914, *A. medemi* Gans & Mathers, 1977, *A. metallurga*, *A. mongoyo*, *A. munoai* Klappenbach, 1960, *A. myersi* Hoogmoed, 1989, *A. nana* Perez & Borges-Martins, 2019, *A. nigricauda* Gans, 1966, *A. occidentalis* Cope, 1875, *A. pericensis* Noble, 1921, *A. plumbea* Gray, 1872, *A. polygrammica* Werner, 1901, *A. prunicolor* (Cope, 1885), *A. ridleyi* Boulenger, 1890, *A. rozei* Lancini, 1963, *A. sanctaeritae* Vanzolini, 1994, *A. saxosa* (Castro-Mello, 2003), *A. schmidt* Gans, 1964, *A. slateri* Boulenger, 1907, *A. slevini* Schmidt, 1936, *A. spurrelli* Boulenger, 1915, *A. steindachneri* Strauch, 1881, *A. supernumeraria* Mott, Rodrigues & Dos Santos, 2009, *A. talisiae* Vanzolini, 1995, *A. tyaraju*, Perez & Borges-Martins, 2019, *A. townsendi* Stejneger, 1911, *A. tragorhectes* Vanzolini, 1971, *A. uroxena*, *A. vanzolinii* Gans, 1963, *A. vermicularis*, and *A. xera* Thomas, 1966.

Amphisbaena amethysta sp. nov. differs from *Amphisbaena* species with four precloacal pores mainly by following combination of meristic characters (Table 1): cephalic shield distinct (vs frontals and parietals shields not distinct in *A. supernumeraria*, ocular and second supralabial not distinct in *A. cubana*); snout slightly compressed (vs hard compressed in *A. kingi* and rounded in all other species, except *A. kraoh*, *A. brasiliiana*, *A. bahiana*, *A. bedai*, and *A. saxosa*); 185–199 dorsal half-annuli (vs < 170 annuli in *A. cayemite* and > 200 annuli in *A. acangaoba*, *A. arda*, *A. arenaria*, *A. bahiana*, *A. bakeri*, *A. barbouri*, *A. bedai*, *A. borellii*, *A. brasiliiana*, *A. caeca*, *A. carlgansi*, *A. carvalhoi*, *A. cayemite*, *A. cuiabana*, *A. cunhai*, *A. elbakyanae*, *A. fenestrata*, *A. frontalis*, *A. gonavensis*, *A. gracilis*, *A. hastata*, *A. hoogmoedi*, *A. kingi*, *A. kraoh*, *A. lumbricalis*, *A. manni*, *A. medemi*, *A. mongoyo*, *A. munoai*, *A. myersi*, *A. nigricauda*, *A. occidentalis*, *A. plumbea*, *A. polygrammica*, *A. rozei*, *A. sanctaeritae*, *A. saxosa*, *A. slevini*, *A. spurrelli*, *A. steindachneri*, *A. supernumeraria*, *A. talisiae*, *A. tyaraju*, *A. townsendi*, *A. uroxena*, *A. vanzolinii*, *A. vermicularis*, and *A. xera*); 13–16 caudal annuli (vs > 16 annuli in *A. albocingulata*, *A. arda*, *A. arenaria*, *A. arenicola*, *A. bedai*, *A. bolivica*, *A. borellii*, *A. carvalhoi*, *A. caudalis*, *A. cegei*, *A. cuiabana*, *A. cunhai*, *A. darwini*, *A. elbakyanae*, *A. frontalis*, *A. gracilis*, *A. hastata*, *A. hoogmoedi*, *A. leali*, *A. lumbricalis*, *A. manni*, *A. medemi*, *A. metallurga*, *A. mongoyo*, *A. munoai*, *A. myersi*, *A. nana*, *A. nigricauda*, *A. occidentalis*, *A. polygrammica*, *A. prunicolor*, *A. rozei*, *A. sanctaeritae*, *A. saxosa*, *A. schmidt*, *A. slateri*, *A. slevini*, *A. spurrelli*, *A. steindachneri*, *A. supernumeraria*, *A. talisiae*, *A. tyaraju*, *A. townsendi*, *A. tragorhectes*, and *A. vermicularis*); conspicuous autotomic site between 4th–6th caudal annuli (vs absent in *A. acangaoba*, *A. alba*, *A. angustifrons*, *A. bakeri*, *A. barbouri*, *A. brasiliiana*, *A. carioca*, *A. carlgansi*, *A. cayemite*, *A. cunhai*, *A. fenestrata*, *A. gonavensis*, *A. hastata*, *A. hoogmoedi*, *A. innocens*, *A. lumbricalis*, *A. occidentalis*, *A. ridleyi*, *A. saxosa*, and *A. uroxena*; or from the 7th caudal annuli in *A. albocingulata*, *A. arda*, *A. arenicola*, *A. carvalhoi*, *A. cegei*, *A. cuiabana*, *A. darwini*, *A. heathi*, *A. kingi*, *A. metallurga*, *A. mongoyo*, *A. myersi*, *A. nana*, *A. prunicolor*, *A. schmidt*, *A. slateri*, *A. steindachneri*, *A. supernumeraria*, *A. talisiae*, *A. tyaraju*, *A. townsendi*, *A. tragorhectes*, and *A. vanzolinii*), smooth and rounded tail tip




Figure 4. Variation (grey in the drawings) of cephalic shields of *Amphisbaena amethysta* sp. nov. **A** dorsal view of the head of CEPB 2280 with frontal variation **B** dorsal view of CEPB 2309 with four parietal shields, and **C** lateral view of left side of CEPB 2303 (paratype) with supralabials fused. Scale bar: 3 mm.

[vs bluntly ridged tail tip in *A. bahiana*; slightly dorsally compressed in *A. acan-*
gaoba; with tubercles sit is depressed (compressed dorsoventrally) in *A. urox-*
ena; with modified conic pointed tubercles in *A. caetitensis*; and vertically keeled

in *A. borellii* and *A. steindachneri*]; 16–21 dorsal segments at midbody (vs < 16 in *A. albocingulata*, *A. arenaria*, *A. arenicola*, *A. barbouri*, *A. carlgansi*, *A. carvalhoi*, *A. cayemite*, *A. cuiabana*, *A. elbakyanae*, *A. fenestrata*, *A. heathi*, *A. hogei*, *A. elbakyanae*, *A. metallurga*, *A. munoai*, *A. nana*, *A. nigricauda*, *A. sanctaeritae*, *A. schmidtii*, *A. slateri*, *A. slevini*, *A. supernumeraria*, *A. talisiae*, *A. tyaraju*, *A. tragorrrhectes*, *A. uroxena*, and *A. vanzolinii*); and > 21 segments in *A. alba*, *A. arda*, *A. bolivica*, *A. camura*, *A. hoogmoedi*, and *A. kraoh*); 16–21 ventral segments at midbody (vs < 16 in *A. rozei*, *A. sanctaeritae*, and *A. tragorrrhectes*; and > 21 segments in *A. alba*, *A. arda*, *A. bolivica*, *A. camura*, *A. cegei*, *A. gonavensis*, *A. hyporissor*, *A. kraoh*, *A. occidentalis*, and *A. townsendi*); 3/3 supralabials (vs 2/2 in *A. slevini* and *A. vanzolinii*; and 4/4 in *A. acangaoba*, *A. alba*, *A. angustifrons*, *A. arda*, *A. arenaria*, *A. bedai*, *A. camura*, *A. cayemite*, *A. occidentalis*, *A. plumbea*, *A. ridleyi*, *A. saxosa*, *A. townsendi*, *A. tragorrrhectes*, and *A. vermicularis*); and 3/3 infralabials (vs 2/2 in *A. slevini* and *A. vanzolinii*; and or 4/4 in *A. occidentalis*, *A. plumbea*, *A. ridleyi*, *A. saxosa*, *A. townsendi*, and *A. tragorrrhectes*).

Table 1. Diagnostic characters for the *Amphisbaena* species with four precloacal pores. PC – precloacal pore, CS – cephalic shield, DA – dorsal half-annulus, CA – caudal annulus, AS – autotomic site, DS – dorsal segment, VS – ventral segment, SL – supralabial, IL – infralabial, PM – postmalar, SN – snout, DG – dorsal groove, VG – ventral groove, a – absent, cp – compressed, n/a – non-available data, p – present, rd – rounded, and sc – slightly compressed. Differences from the new species in bold.

Species	PC	CS	DA	CA	AS	DS	VS	SL	IL	PM	SN	DG	VG	Reference
<i>A. amethysta</i> sp. nov.	4	distinct and paired	185–199	13–16	4–6	16–21	16–21	3	3	a	sc	a	a	present study
<i>A. acangaoba</i>	4–8	distinct and paired	216–293	13–17	a	18–24	18–24	4	3	p	rd	a	a	Ribeiro et al. (2020)
<i>A. alba</i>	4–10	distinct and paired	198–248	13–21	a	30–42	35–46	4	3	p	rd	a	a	Gans (1962a)
<i>A. albocingulata</i>	4	distinct and paired	190–204	24–27	8–9	12–14	16–18	3	3	a	rd	a	a	Perez et al. (2012)
<i>A. angustifrons</i>	3–6	distinct and paired	190–215	12–18	a	20–31	21–30	4	3	p	rd	a	a	Gans (1965a); Gans and Diefenbach (1972)
<i>A. arda</i>	4	distinct and paired	242	30	8	23	23	4	3	p	rd	a	a	Rodrigues et al. (2003)
<i>A. arenaria</i>	4	distinct and paired	285–307	22–23	6–7	12–14	14–16	4	3	a	rd	a	a	Teixeira Junior et al. (2016)
<i>A. arenicola</i>	4	distinct and paired	199–216	20–22	8–9	12–14	16–18	3	3	P	rd	n/a	n/a	Perez and Borges-Martins (2019)
<i>A. bahiana</i>	4	distinct and paired	204–223	14–16	4–5	12–16	14–16	3	3	p	sc	p	p	Gans (1964b); Dal Vechio et al. (2018)
<i>A. bakeri</i>	4	distinct and paired	239–255	14–16	a	14–16	16–17	3	3	a	rd	n/a	n/a	Gans and Alexander (1962)
<i>A. barbouri</i>	4–6	distinct and paired	226–240	13–18	a	12–14	16–18	3	3	a	rd	n/a	n/a	Gans and Alexander (1962); Thomas (1966)
<i>A. bedai</i>	4	distinct and paired	272–284	22–23	6	18–20	16–18	4	3	p	sc	a	a	Oliveira et al. (2018)
<i>A. bolivica</i>	4–6	distinct and paired	200–231	18–26	4–5	27–36	26–36	3	3	p	rd	a	a	Montero (1996)
<i>A. borellii</i>	4	distinct and paired	239–245	17–19	6–8	14–16	16–20	3	3	n/a	rd	a	a	Gans (1964b) and Oliveira et al. (2018)
<i>A. brasiliiana</i>	4	distinct and paired	213–229	11–15	a	18–21	18–22	3	3	p	sc	P	a	Oliveira et al. (2018)
<i>A. caeca</i>	4–6	distinct and paired	217–236	13–18	4–8	13–18	14–20	3	3	p	rd	n/a	n/a	Gans and Alexander (1962)
<i>A. caetitensis</i>	4	distinct and paired	186–194	10–12	a	16	19–22	3	3	a	rd	a	a	Almeida et al. (2018)
<i>A. camura</i>	4–6	distinct and paired	188–207	14–19	4–5	28–42	29–46	4	3	p	rd	a	a	Gans (1965a); Hoogmoed and Ávila-Pires (1991)

Species	PC	CS	DA	CA	AS	DS	VS	SL	IL	PM	SN	DG	VG	Reference
<i>A. carioca</i>	4	distinct and paired	186	21	a	10–12	n/a	3	3	p	rd	a	a	Rocha et al. (2023)
<i>A. carlgansi</i>	4	distinct and paired	212–228	14–16	a	14	18–20	3	3	a	rd	a	a	Thomas and Hedges (1998)
<i>A. carvalhoi</i>	4	distinct and paired	231–245	19–22	7–8	12–14	16–18	3	3	a	rd	a	a	Gans (1965b)
<i>A. caudalis</i>	4	distinct and paired	193–217	17–21	6	12–16	18–21	3	3	a	rd	n/a	n/a	Gans and Alexander (1962); Thomas and Hedges (1998)
<i>A. cayemite</i>	4	distinct and paired	150–164	10–13	a	12–13	18	4	3	p	rd	n/a	n/a	Thomas and Hedges (2006)
<i>A. cegei</i>	4	distinct and paired	198	22	7	21–22	22–23	3	3	a	rd	n/a	n/a	Montero et al. (1997)
<i>A. cubana</i>	4–6	ocular and second SL fused	199–218	10–16	6–9	12–16	14–18	3	3	a	rd	n/a	n/a	Gans and Alexander (1962); Thomas (1966)
<i>A. cuiabana</i>	4	distinct and paired	286–292	18–20	9–10	14	16	3	3	a	rd	p	a	Strüssmann and Carvalho (2001)
<i>A. cunhai</i>	4	distinct and paired	226–239	25–26	a	14–16	14–18	3	3	p	rd	a	a	Hoogmoed and Ávila-Pires (1991)
<i>A. darwini</i>	2–5	distinct and paired	174–195	19–25	7–10	13–19	16–23	3	3	n/a	rd	a	a	Perez et al. (2012)
<i>A. elbakyanae</i>	4	distinct and paired	245–257	20–24	6–7	13–15	16–18	3	3	p	rd	p	a	Torres-Ramírez et al. (2022)
<i>A. fenestrata</i>	4	distinct and paired	236–249	12–14	a	13–14	14–16	3	3	p	rd	n/a	n/a	Gans and Alexander (1962)
<i>A. frontalis</i>	0–4	distinct and paired	235–275	23–29	5–7	14–18	14–16	3	3	a	rd	a	a	Ribeiro-Júnior et al. (2022)
<i>A. gonavensis</i>	4	distinct and paired	214–225	10–13	a	15–18	22–25	3	3	p	rd	n/a	n/a	Gans and Alexander (1962)
<i>A. gracilis</i>	4	distinct and paired	224–248	21–24	6–7	13–16	14–17	3	3	p	rd	p	a	Gonzales-Sponga and Gans (1971)
<i>A. hastata</i>	4	distinct and paired	266–273	40	a	18	16	3	3	a	rd	a	p	Vanzolini (1991)
<i>A. heathi</i>	4	distinct and paired	183–187	n/a	7–8	12	18–20	3	3	a	rd	a	a	Gans (1965b)
<i>A. hogei</i>	4	distinct and paired	177–191	15–19	4–7	10–13	14–18	3	3	p	rd	a	a	Gans (1966); Vanzolini (1950)
<i>A. hoogmoedi</i>	4	distinct and paired	247–252	27	7–8	22–24	19–21	3	3	a	sc	a	a	Oliveira et al. (2018)
<i>A. hyporissor</i>	4	distinct and paired	199–227	16–21	5	14–18	22–24	3	3	p	rd	n/a	n/a	Thomas (1965); Thomas and Hedges (2006)
<i>A. innocens</i>	4	distinct and paired	185–220	11–14	a	13–16	18–21	3	3	a	rd	n/a	n/a	Gans and Alexander (1962); Thomas and Hedges (1998)
<i>A. kingi</i>	4	distinct and paired	214–244	15–23	7	12–19	14–22	3	3	p	cp	a	a	Gans and Rhodes (1964); Oliveira et al. (2018)
<i>A. kraoh</i>	4–6	distinct and paired	270–281	15	5	28	24–27	3	3	n/a	sc	n/a	n/a	Oliveira et al. (2018)
<i>A. leali</i>	4	distinct and paired	188–206	17–20	6	14–16	20–22	3	3	a	rd	n/a	n/a	Thomas and Hedges (2006)
<i>A. lumbricalis</i>	2–6	distinct and paired	225–247	20–26	a	12–16	16–20	3	3	n/a	rd	a	a	Vanzolini (1996)
<i>A. manni</i>	4–9	distinct and paired	209–243	17–22	5–7	12–16	14–20	3	3	a	rd	n/a	n/a	Gans and Alexander (1962)
<i>A. medemi</i>	4	distinct and paired	230–235	17–18	5–7	16	18	3	3	a	rd	a	p	Gans and Mather (1977)
<i>A. metallurga</i>	2–4	distinct and paired	185–199	23–25	7–9	12–14	14–16	3	3	p or a	rd	a	a	Costa et al. (2019)
<i>A. mongoyo</i>	4	distinct and paired	208	25	10	14	16	3	3	a	rd	a	a	Teixeira Junior et al. (2019)
<i>A. munoai</i>	4	distinct and paired	202–218	18–23	5–9	10–14	13–18	3	3	p	rd	a	a	Perez and Borges-Martins (2019)
<i>A. myersi</i>	4	distinct and paired	221	28	8	16	16	3	3	n/a	rd	a	a	Hoogmoed (1989)
<i>A. nana</i>	4	distinct and paired	195–216	18–22	7–10	12–14	14–17	3	3	p	rd	n/a	n/a	Perez and Borges-Martins (2019)
<i>A. nigricauda</i>	4	distinct and paired	222–226	19–24	6–9	10	16	3	3	a	rd	a	a	Gans (1966)
<i>A. occidentalis</i>	4	distinct and paired	261–279	18–21	a	16–19	22–27	4	4	p	rd	p	p	Gans (1961)

Species	PC	CS	DA	CA	AS	DS	VS	SL	IL	PM	SN	DG	VG	Reference
<i>A. pericensis</i>	4	distinct and paired	198–218	16–19	6–8	12–16	16–20	3	3	a	rd	a	a	Gans (1963a)
<i>A. plumbea</i>	4	distinct and paired	233–282	16–21	5–9	18–27	20–30	4	4	p	rd	a	a	Gans and Diefenbach (1972)
<i>A. polygrammica</i>	4	distinct and paired	270	20	n/a	18	16	n/a	n/a	n/a	n/a	n/a	n/a	Vanzolini (2002)
<i>A. prunicolor</i>	4	distinct and paired	181–215	18–27	7–10	10–17	14–20	3	3	p	rd	a	a	Perez et al. (2012)
<i>A. ridleyi</i>	4	distinct and paired	172–192	14–17	a	16–18	20–28	4	4	p	rd	a	a	Gans (1963b)
<i>A. rozei</i>	4	distinct and paired	205–209	20	6 or a	15–16	14	3	3	p	rd	a	a	Vanzolini (2002); Costa et al. (2019)
<i>A. sanctaeritae</i>	4	distinct and paired	269–288	18–20	6–7	10	14	3	3	p	rd	a	a	Costa et al. (2019)
<i>A. saxosa</i>	4	distinct and paired	253–272	17–21	a	18–24	16–21	4	4	p	sc	a	a	Oliveira et al. (2018)
<i>A. schmidtii</i>	4	distinct and paired	198–202	20–22	7–8	14	16–17	3	3	p	rd	a	a	Gans (1964a)
<i>A. slateri</i>	4	distinct and paired	176–213	20–24	7–10	10–14	14–16	3	3	p or a	rd	a	a	Gans (1967); Costa et al. (2018)
<i>A. slevini</i>	4	distinct and paired	204–211	23–25	5–6	10–14	10–12	2	2	a	rd	p	a	present study
<i>A. spurrelli</i>	4	distinct and paired	218–222	18–20	6–7	16–18	16–18	3	3	p	rd	a	a	Gans (1962b); Costa (2020)
<i>A. steindachneri</i>	4	distinct and paired	256–266	17–18	7	14–16	16	3	3	a	rd	p	p	Gans (1964b)
<i>A. supernumeraria</i>	4	not distinct	333–337	22–23	10–12	14	17–18	3	3	n/a	rd	a	n/a	Mott et al. (2009)
<i>A. talisiae</i>	4	distinct and paired	205–234	17–29	7	10–14	14–18	3	3	a	rd	a	a	Vanzolini (1995); Costa et al. (2019)
<i>A. tyaraju</i>	4	distinct and paired	204–221	18–25	7–9	10–14	13–16	3	3	p	rd	n/a	n/a	Perez and Borges-Martins (2019)
<i>A. townsendi</i>	4	distinct and paired	261–279	22–26	7–8	16–19	22–27	4	4	p	rd	n/a	n/a	Gans (1961)
<i>A. tragorhectes</i>	4	distinct and paired	196	31	12	12	12	4	4	p	rd	a	a	present study
<i>A. uroxena</i>	0–4	distinct and paired	210–213	12–13	a	14	14–15	3	3	p	rd	a	p	Mott et al. (2008)
<i>A. vanzolinii</i>	4	distinct and paired	225–228	n/a	7	12–13	16–17	2	2	p	rd	a	a	Gans (1963c)
<i>A. vermicularis</i>	4	distinct and paired	211–254	23–34	5–7	18–26	18–25	4	3	n/a	rd	a	a	Gans and Amdur (1966)
<i>A. xera</i>	4	distinct and paired	225–234	13–16	5	12–16	14–16	3	3	a	rd	a	a	Thomas (1966)

Description of the holotype. medium-sized specimen; snout-vent length 233 mm plus 0.50 mm of cloacal portion; tail length 21.24 mm, representing 9.1% of snout-vent length; midbody diameter 8.2 mm (3.5% of snout-vent length); head relatively small, 6.90 mm (~ 2.9% of snout-vent length); snout compressed in dorsal view and slightly convex in profile view, hardly keratinised, rostrum projecting forward beyond the jaw (prognathous). Anterior portion of body is slightly narrower. Rostral subtriangular, visible in dorsal and ventral view (Fig. 1), almost as high (2.21 mm) as wide (2.11 mm), in contact with nasal and first supralabial lateroposteriorly. Nasals subrectangular, aligned at the midline (1.00 mm suture) (Fig. 1A), almost as long (2.05 mm) as wide (1.99 mm), in contact with first supralabial laterally and prefrontals posteriorly, with nostrils placed near the antero-inferior angle of the nasal shield (Fig. 1B).

Prefrontals paired, relatively large (41.6% of head length), with a shorter middorsal suture (2.01 mm; 29.3% of head length), longer than the nasal middorsal suture (1.00 mm, 14.6% of head length), almost as long as frontal middorsal suture (2.10 mm, 30.6% of head length), anterior border convex, lateroposterior portion projected, in contact with second supralabial and ocular laterally, frontals posteriorly and in point contact with postocular posteriorly

(Fig. 1A). Frontals subtriangular, longer (suture length) than wide (1.58 mm), aligned at midline (2.10 mm), in narrow contact with the oculars, and in broad contact posterolaterally with the postocular and parietal. Parietals in two larger irregular segments, wider (1.48 mm) than long (1.04 mm), intercalated by four very small segments; not aligned at the midline, in narrow contact with postoculars laterally, and followed by the first dorsal half-annulus. Occipitals absent (Fig. 1A). Oculars almost diamond-shape, almost as long (1.57 mm) as high (1.51 mm), in contact with prefrontals and second supralabial anteriorly, third supralabial and postocular posteriorly, and in point contact with postsupralabial. Eyes slightly visible. Postocular longer than wide, sub-pentagonal, in contact with frontal, labial, parietal and the segments of the first dorsal half-annulus, and in point contact with prefrontal. Temporal subrectangular, higher (1.55 mm) than long (0.92 mm), in contact with third supralabial anteriorly, postocular and postsupralabial laterally and first dorsal half-annulus posteriorly (Fig. 1B).

Three supralabials, irregularly polygonal; first subtrapezoid, longer (2.05 mm) than high (1.50 mm), in contact with second supralabial posteriorly; second supralabial sub-pentagonal, higher (1.76 mm) than long (1.59 mm maximum length), in contact with prefrontal, ocular and third supralabial; third supralabial subtrapezoid, almost as high (1.37 mm) as long (1.24 mm), in contact with ocular and postsupralabial. Postsupralabial subquadrangular, representing almost 50% of third supralabial high, in contact with temporal laterally and first half-annulus posteriorly (Fig. 1B). Mental longer (1.72 mm) than wide, anterior border wider (1.72 mm) than posterior (1.08 mm), in contact with the first pair of infralabials and postmental. Postmental longer (2.01 mm) than wide (1.65 mm), in contact with the first and second infralabial, narrowly with malar, and two anterior postgenials. Postgenials with five shields irregularly distributed, in contact with malars and first ventral half-annuli (Fig. 1C).

Three infralabials, first medium sized, irregular polygonal, almost as long (1.55 mm) as wide (1.56 mm), in contact with second supralabial; second the largest, sub-pentagonal, wider (2.36 mm) than long (1.81 mm maximum length), in contact with third infralabial; third infralabial smallest, almost as long (1.24 mm) as wide (1.37 mm) (Fig. 1C).

Body annuli well demarcated, first and second annuli without enlarged dorsal segments. Segments become regularly rectangular toward posterior portion of body and progressively longer than wide, and smaller in size, and larger towards midventral areas starting from the fifth half-annulus. One hundred ninety-four dorsal and 195 ventral half-annuli, three lateral half-annuli, 14 caudal annuli plus tip rounded; tail relatively long with autotomy line on the fifth annulus, 18/18 dorsal and ventral segments at midbody, respectively and 28 segments in fourth caudal annulus. Lateral sulci clearly visible from the forty-ninth annulus; dorsal and ventral sulci absent. Cloacal plate with six segments increasing in size from towards midline, eleven postcloacal segments; four precloacal pores strongly visible on the row of segments on the last ventral half-annulus; each pore placed on the posterior half of a single segment, and distributed along a continuous series of segments, but pores in the medial scales placed laterally (Fig. 2).

Intraspecific variation. The main variations in the type series for meristic and morphometric data are given in Table 2. Variation in the arrangement and contact of shields were also observed. CEPB 2280 presents frontal fused with the parietal and segments of first and second body annuli (Fig. 3A), CEPB 2309

Table 2. Variation in meristic and morphometric (mm) data in the type series of *Amphisbaena amethysta* sp. nov. S – sex, DA – dorsal half–annuli, LA – lateral half–annuli adjacent to cloacal region, VA – ventral half–annuli, CA – caudal annuli, AS – autotomic site, DS – dorsal segment in midbody, VS – ventral segment in midbody, SCA – segments of fourth caudal annulus, PRCL – precloacal segments, POCL – postcloacal segments, PGE – postgenials, PA – parietals, SVL – snout–vent length, TL – tail length, BW – body width in midbody, bs – brooked specimen, Ita – last tail annuli, n/a – non-available data, +n – mutilated tail, * – tail cicatrised, ** – tail not cicatrised, un- unidentified.

Specimens	S	DA	LA	VA	CA	AS	DS	VS	SCA	PRCL	POCL	PGE1	PA	SVL	TL	BW
CEPB 2311 (Holotype)	male	194	3	195	14	5	18	18	28	6	11	5	1/1	233	21.2	8.2
CEPB 2301 (Paratype)	female	191	3	191	5 + n	Ita5**	19	17	29	7	13	5	1/1	206	3.5 + n	6.4
CEPB 2302 (Paratype)	male	194	4	192	15	5	20	18	31	7	13	5	1/1	199	17.8	7.2
CEPB 2303 (Paratype)	female	194	4	197	5 + n	Ita5*	17	16	27	6	14	5	1/1	197	8.2 + n	7.0
CEPB 2308 (Paratype)	female	195	4	196	15	5	18	18	28	n/a	13	4	1/1	150	13.0	5.1
CEPB 2327 (Paratype)	female	196	3	196	14	5	20	19	31	6	13	5	1/1	205	16.5	6.0
CEPB 2331 (Paratype)	female	197	4	195	15	5	18	16	30	6	11	5	1/1	175	14.4	5.8
CEPB 2346 (Paratype)	male	195	4	195	14	5	18	18	29	6	14	5	1/1	138	11.3	4.3
CEPB 2379 (Paratype)	male	185	4	189	5 + n	Ita5**	19	18	29	6	13	5	1/1	180	5.0 + n	7.0
CEPB 2381 (Paratype)	female	189	4	189	15	4	16	16	24	6	12	5	1/1	165	14.8	5.6
CEPB 2298	female	199	3	199	14	n/a	18	18	30	7	13	5	1/1	215	16.7	7.4
CEPB 2299	female	193	4	193	6+n	Ita5*	18	15	29	6	13	4	1/1	186	7.6 + n	5.1
CEPB 2300	male	194	3	195	14	5	18	17	26	6	12	5	1/1	213	15.3	6.1
CEPB 2304	female	190	4	192	14	5	18	n/a	26	6	12	5	1/1	155	11.6	3.5
CEPB 2305	female	192	4	192	15	5	18	16	28	6	13	6	1/1	170	14.8	5.6
CEPB 2306	female	197	4	196	14	4	18	18	28	8	12	5	1/1	145	10.8	4.2
CEPB 2307	male	190	3	190	6 + n	Ita6**	18	17	27	6	13	5	2/2	220	6.8 + n	7.9
CEPB 2309	un	196	3	197	6 + n	Ita5*	19	16	n/a	6	14	6	1/2	125	7.9 + n	4.5
CEPB 2310	female	193	4	194	14	5	18	18	26	6	11	2	0/1	195	15.7	5.5
CEPB 2312	female	192	3	192	15	5	18	18	30	6	15	3	1/1	190	15.1	6.6
CEPB 2313	female	189	3	190	15	5	18	15	28	6	14	5	1/1	130	11.7	4.2
CEPB 2314	female	191	3	193	14	5	18	16	26	6	13	6	1/1	185	11.9	5.7
CEPB 2315	female	193	3	193	14	5	18	19	27	6	12	5	1/1	177	14.1	5.9
CEPB 2316	female	195	3	195	16	6	20	18	28	7	13	5	1/1	173	15.2	6.4
CEPB 2317	female	195	5	196	5 + n	Ita5**	18	n/a	n/a	6	13	5	1/1	180	3.0 + n	5.3
CEPB 2318	female	196	3	195	15	5	20	18	31	6	11	5	1/1	120	16.1	4.3
CEPB 2319	female	186	4	186	5 + n	Ita5**	19	18	29	6	12	5	1/1	95	2.0 + n	3.2
CEPB 2320	female	185	4	182	15	5	19	18	26	7	13	2	1/1	92	7.7	2.6
CEPB 2321	female	194	3	192	16	6	18	18	30	6	12	5	1/1	160	14.1	6.0
CEPB 2322	female	196	4	195	5 + n	Ita5**	21	18	27	6	11	4	1/1	155	3.7 + n	5.3
CEPB 2323	female	194	4	193	15	5	16	16	24	6	13	2	1/1	112	8.8	3.0
CEPB 2324	male	193	3	193	16	6	19	18	31	6	13	5	1/1	162	14.6	5.7
CEPB 2325	female	193	4	190	14	5	19	18	27	6	11	5	1/1	174	14.2	6.1
CEPB 2326	male	198	3	197	16	6	17	17	28	6	10	5	1/1	195	16.4	6.4
CEPB 2328	female	192	3	194	15	6	18	18	28	6	12	5	1/1	173	15.5	6.5
CEPB 2329	male	195	4	196	15	5	19	18	30	6	13	6	1/1	170	13.0	5.7
CEPB 2330	male	194	4	193	5 + n	Ita5**	20	18	30	6	13	5	1/1	178	2.6 + n	6.0
CEPB 2332	male	190 + n	4	190	15	5	18	16	28	6	15	5	1/1	bs	11.1	4.0
CEPB 2333	female	196	3	194	15	n/a	18	17	26	6	10	2	1/1	203	16.9	6.3
CEPB 2334	female	194	3	192	5 + n	5	20	18	30	6	14	6	1/1	190	7.0 + n	7.3
CEPB 2335	female	197	3	197	14	5	20	18	31	6	12	5	1/1	135	9.7	3.9
CEPB 2336	female	190	4	190	15	6	18	17	29	6	13	4	1/1	105	9.1	3.4
CEPB 2337	female	186	4	185	5 + n	Ita5**	19	18	29	6	13	5	1/1	145	3.7 + n	4.9
CEPB 2338	female	190 + n	4	186	13	5	18	17	28	6	12	5	1/1	~ 160	13.8	5.1
CEPB 2339	female	192	4	193	16	5	19	18	32	6	13	5	1/1	127	10.6	4.0
CEPB 2356	male	188	3	189	16	n/a	18	18	28	7	13	3	1/1	190	16.7	5.8
CEPB 2380	female	192	3	195	14	5	17	16	25	6	12	5	2/2	140	11.5	5.6
Minimum		185	3	182	13	4	16	15	24	6	10	2		92	2.2	2.6
Maximum		199	5	199	16	6	21	19	32	8	15	6	n/a	233	21.2	8.2
Mean		192.9	3.6	193.0	14.8	5.0	18.4	17.4	28.3	6.2	12.6	4.7	n/a	167.3	13.4	5.5
Mode		194	4	195	15	5	18	18	28	6	13	5	n/a	190	n/a	n/a
Standard deviation		3.4	0.5	3.4	0.8	0.4	1.0	1.0	1.9	0.5	1.1	1.0	n/a	34.1	3.5	1.3

present four parietals shields (Fig. 3B), and CEPB 2303 present supralabials fused in left side (Fig. 3C).

Colour in life. Dorsum and lateral parts with dark brown coloration on the segments, which is more pronounced in the vertebral (Fig. 4A) and dorsal section of the tail regions (Fig. 4B). Pink predominates in areas where the brown colour is less pronounced. We do not have photographs of the ventral region of the live specimen.

Colour in preservative (ethylic alcohol 70%). Dorsum cream, with brown colouring on the segments in the dorsum, lateral parts, and dorsal portions of the tail portions. Dorsal, lateral, and ventral portions of head pale brown, darker than the ventral portion. Venter cream coloured.

Etymology. The specific epithet refers to the mineral amethyst that is a type of quartz and also the name of the region of the type locality “Brejinho das Ametistas”, a district located south of the municipality of Caetité, state of Bahia. This region has been an amethyst mining centre since the beginning of the 20th century. Spix and Martius (1938) defined the mineral from the “Brejinho das Ametistas” mines as “the beautiful amethysts” on their trip through the “Alto Sertão” of Bahia at the beginning of the 19th century (Cotrim 2015). The region currently has an economy based on mining and energy activities focused on wind energy production. The type series was collected during the execution of environmental programs within the scope of Bahia Mineração (BAMIN), which operates in the exploration of iron ore in the “Brejinho das Ametistas” region.

Distribution and habitat. *Amphisbaena amethysta* sp. nov. is known from municipality of Caetité municipality, state of Bahia, Brazil (Fig. 5).




Figure 5. Geographic distribution of *Amphisbaena amethysta* sp. nov. and others *Amphisbaena* species from Espinhaço Mountain Range locality. Black arrows indicate the type localities of the species with more than one known geographic record.

The region is in the northern portion of the Espinhaço Mountain Range, has an average altitude of 1000 m a.s.l., and lies within the ecotone between two morphoclimatic domains, Caatinga and Cerrado. In the region there are patches of deciduous and semi-deciduous forests [“Floresta Estacional Decidual” and “Floresta Semidecidual Montana” sensu IBGE (2023)] associated with valleys, slopes, and gallery forests, and containing floristic elements common to the vegetation of the Caatinga, Cerrado, and Atlantic Forest morphoclimatic domains. Areas of savannah vegetation with rock outcrops, typical of the woody Caatinga, occur at higher elevations (Fig. 6).

Phylogenetic relationships. Our concatenated alignment totalled 4806 base pairs (1007 bp for 12S, 528 bp for 16S, 761 bp for *nd2*, 679 bp for *bdnf*, 574 bp for *c-mos*, and 1257 bp for *rag1*). Partition Finder identified a best-fit scheme composed of ten partitions with the GTR + G model. The resulting ML topology (Fig. 7) for the higher-level affinities was similar to those reported by previous studies





Figure 6. Caatinga site where the holotype of *Amphisbaena amethysta* sp. nov. was collected in the northern portion of Espinhaço Mountain Range, in state of Bahia, Brazil.



(Mott and Vieites 2009; Longrich et al. 2015; Graboski et al. 2022) (Fig. 7). *Amphisbaena amethysta* sp. nov. was recovered as a sister group of *A. caetitensis*, with 92% of bootstrap support. The clade composed by *Amphisbaena amethysta* sp. nov. and *A. caetitensis* was recovered as a sister group of *A. angustifrons*, *A. darwini*, *A. kingi*, *A. leeseri*, and *A. munoai* with low bootstrap support (21%). The genetic distance (*p*-distance) between *Amphisbaena amethysta* sp. nov. and *A. caetitensis* is 6.1% for 16S.

Discussion

Amphisbaena amethysta sp. nov. is the 73rd species of the genus with four precloacal pores, the 22nd species from Caatinga, and sixth species with a restricted distribution from this morphoclimatic domain. The recognition of *A. amethysta* as a new species is based on molecular data and on set of morphological characters that includes four precloacal pores, a slightly compressed snout, 185–199 dorsal and 182–199 ventral half-annuli, 13–16 caudal annuli, 16–21 dorsal and ventral segments in the midbody, 3/3 supralabials and infralabials, a smooth and rounded tail tip, an autotomic site between 4th and 6th caudal annuli, and by the absence of fusion of the cephalic shields, postmalar shields, and of dorsal and ventral grooves.

The new species is most closely related to *A. caetitensis* (from an elevation of 854 m in the municipality of Caetité, state of Bahia), a sister species with a genetic distance (*p* distance) of 6.1% for the 16S gene, and which differs morphologically mainly in the modified conic pointed tubercles on the tip tail and absence of an autotomic site (see diagnosis). This tail shape appears to have arisen independently in *A. caetitensis* and *A. uroxena*. Almeida et al. (2018) reported a genetic distance of the 7.65% between *A. uroxena* and *A. caetitensis*, and stated that Bayesian inference does not allow a clear resolution of the relationship between the two species. Our phylogenetic analyses corroborate the results of Almeida et al. (2018), and do not structure a clade for species with the modified conic pointed tubercles on the tip tail. We found an even lower distance between *A. caetitensis* and *A. amethysta* sp. nov. and a phylogenetic correlation between the species (Fig. 7), and recovered *A. uroxena* with sister species of *A. mongoyo*, structuring a clade with *A. longinqua* and *A. bahiana* (Clade G05, Fig. 7). Additionally, with a low support, we retrieved the clade formed by *Amphisbaena amethysta* sp. nov. and *A. caetitensis* as a sister group to clade G02 (Fig. 7) formed by *A. leeseri*, *A. darwini*, *A. munoai*, *A. angustifrons*, and *A. kingi*. This grouping showed a relationship between the species from the Espinhaço Mountain Range and those in other areas of the Atlantic Forest and Cerrado. Additionally, it can indicate a probably convergent evolution between the species with a tuberculate tail shield (*A. caetitensis* and *A. uroxena*) and those species present on mountain within the Espinhaço Mountain Range (clade 5; Fig. 7).

Amphisbaena amethysta appears to be endemic to the northern portion of the Espinhaço Mountain Range with an average altitude of 1000 m a.s.l. Within this, its known extent of occurrence is some 38 km. This distributional pattern is similar to other five species of *Amphisbaena* restricted to the high-altitude areas of the Espinhaço Mountain Range in Bahia [*A. bahiana*, *A. longinqua*, *A. metallurga*, *A. mongoyo*, and *A. uroxena* (Costa et al. 2015; Teixeira Junior et al. 2019)], four of which are closely related phylogenetically, but show no close phylogenetic

relations with *A. amethysta* (*A. metallurga* has no molecular sample available) (Fig. 7). The other three species show no apparent relationship to those from the Espinhaço Mountain Range but also occur at similar altitudes and vegetation types: *A. kiriri* Ribeiro, Gomides & Costa, 2018 (at least 17 km from the nearest *A. bahiana* locality), *A. acangaoba* (occurring in sympatry with *A. kiriri*), and *A. ignatiana* (recorded from the lower sections of the northwestern extremity of the Espinhaço Mountain Range) (Fig. 5) (Roberto et al. 2014; Ribeiro et al. 2018b, 2020).

In the last years morphological characters were commonly used to diagnose *Amphisbaena*, mainly meristics (e.g. Roberto et al. 2014; Costa et al. 2015; Ribeiro et al. 2016, 2019, 2018a, b, 2020; Torres-Ramírez et al. 2021; Ribeiro-Júnior et al. 2022; Rocha et al. 2023) and morphometrics (e.g. Ribeiro et al. 2009; Oliveira et al. 2018) but also with the addition of genetic analyses (e.g., Almeida et al. 2018; Perez and Borges-Martins 2019; Ribeiro et al. 2019), but it is still unclear which characters best represent the interspecific variation of the genus. The identification of diagnostic characters depends on the recognition of intraspecific variations of different species, as well as the analysis of relatively large numbers of samples for each species, the latter being hampered by the fossorial habits of the species in this group and the consequently low frequency of encounters with them. Cryptic species of *Amphisbaena* are rare in bibliographic citations, but recent studies have described new species using diagnoses based on overlapping phenotypic variation and the divergence of the 16S and ND2 genes, complicating species identification using external morphology alone. To reduce the limitations in the identification of new amphisbaenid species, in addition to applying integrative taxonomy whenever possible, we consider it important to standardise the use of meristic and morphometrics characters that do not overlap and to explore new characters, such as the morphometry of the shields (Ribeiro et al. 2008, 2009, 2011, 2015, 2016, 2018a, 2019; Sindaco et al. 2014); analysing characters of internal morphology, such as osteology and gonadal morphology (Ribeiro et al. 2015, 2018a; Oliveira et al. 2018; Angiolani-Larrea et al. 2021); and studying similar species, including type series (Costa et al. 2019; Ribeiro-Júnior et al. 2022).

Amphisbaena amethysta sp. nov. varies in the number of dorsal and ventral half-annuli, caudal annuli, and dorsal and ventral segments in the midbody, the shape and number of parietal shields, and the position of the autotomic site. In addition to variations in shape and number of cephalic shields, specimens of the new species also included individuals with shield fusion, mainly in the parietal region (see discussion of variation). On the other hand, the sample did not vary the number and shape of supra and infralabials and postlabials; the presence of a malar; the shape of the rostral, prefrontal, and frontal shields; the presence and number of precloacal pores; nor in the shape of the segments of the half-annuli of the body and tail. Despite its use as an invariable character, the number of pores can vary intraspecifically. Among the other *Amphisbaena* species with four precloacal pores, 14 vary in the number of precloacal pores: *A. acangaoba*, *A. alba*, *A. angustifrons*, *A. barbouri*, *A. bolivica*, *A. caeca*, *A. camura*, *A. cubana*, *A. darwini*, *A. frontalis*, *A. kraoh*, *A. lumbricalis*, *A. manni*, and *A. uroxena* (Gans 1962a, 1965a; Gans and Alexander 1962; Thomas 1966; Gans and Diefenbach 1972; Hoogmoed and Ávila-Pires 1991; Montero 1996; Vanzolini 1996; Mott et al. 2008; Perez et al. 2012; Ribeiro et al. 2020; Ribeiro-Júnior et al. 2022). Additionally, sexual dimorphism in the presence or absence of pores for *Amphisbaena* has already been reported for *A. frontalis* and *A. uroxena* (Mott et al. 2011; Ribeiro-Júnior et al. 2022).

Mining activities cause several irreversible changes to the environment, including the loss of habitat due to the removal of vegetation and the relocation and excavation of soil during the opening and operation of new mines. Such actions have a direct impact on terrestrial and fossorial species such as amphisbaenians. In this context, applied studies, such as the evaluation of species as biomarkers for metal exposure, mainly with fossorial reptile species, are neglected (Grillitsch and Schiesari 2010; Gil-Jiménez et al. 2021). In such circumstances, the development and execution of environmental programs are critical for effective impact mitigation. Such studies can form valuable tools when conducted by specialised teams with a focus on the scientific use of collected material (Vaz-Silva 2009), and teams concerned with the technical quality of the project, which would result in more careful projects in relation to the sampling design and analytical aspects of the results, allowing greater decision-making power for effective conservation actions (Dias et al. 2019). In this case, environmental studies have made an important contribution to knowledge of the biodiversity of the area under BAMIN's control, Bahia Mineração. Finally, the identification of a new species indicates that the fossorial fauna, as well as that of other groups, in the Espinhaço Mountain Range region is far from being completely known and that it may harbour a much greater diversity of endemic taxa than has been realised so far.

Acknowledgements

We would like to thank the curators and/or collection technicians D. Frost and D. Kizirian (AMNH), G.R. Colli, M. Viana and M. Ayub (CHUNB); S. Carreira (FC-UDELAR); H.M. Chalkidis (LPHA); G. Pontes (MCT-PUCRS); J. Moura-Leite (MHNCI); M. Motte (MNHNP); P. Passo, R. Fernandes and M. Soares (MNRJ); A. Prudente, A. Travassos and F. Sarmento (MPEG); H. Zaher and A. Carvalho (MZUSP); E. Freire and T. Mott (MUFAL); A. Argolo and M. Solé (UESC); V. Ferreira (UFMS/CEUCH/AFB); and F. Curcio (UFMT) for permits, care, and access to specimens. We are grateful to D. M. Soares for photography of the holotype *Amphisbaena amethysta* sp. nov. and to D. Passos, one anonymous reviewer, and the editors N. Yonow and U. Garcia for suggestions that improved the quality of the manuscript. We thank BAMIN Bahia Mineração and the time of Tecsan – Tecnologia e Saneamento responsible for environmental mitigation actions that culminated in the collection of specimens from the type series: Patrícia de Rossi Belufi, Andrezza Sayuri, Victoriano Hirota, Roberto Aires de Assunção, Inácio Fernandes Brito, Fabiana Soares dos Santos, Márcio Santos Borges, and Gabryelle Santos Nascimento. We thanks the National Council for Scientific and Technological Development – CNPq, through the program to support research projects and training human resources for biological taxonomy – PRO-TAX 22/2020, process number 441967/2020-5 CNPq. RG thanks Fundação de Amparo à Pesquisa do Estado de São Paulo for fellowships (FAPESP, 2012/24755-8 and 2016/06866-8) and IGM thanks Fundação de Amparo à Pesquisa do Estado do Pará (FAPESPA, 00000.9.000794/2023).

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding


RG was supported by grants from the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, 2012/ 24755-8 and 2016/06866-8). IGM was supported by grants from the Fundação de Amparo à Pesquisa do Estado do Pará (FAPESPA, 00000.9.000794/2023).

Author contributions

SR, WVS, and APSJ identified the new species. WVS, and RG gathered the molecular data; SR, ECS, APJS, and IGM gathered the morphological data; all authors wrote and revised the text.

Author ORCIDs

Síria Ribeiro  <https://orcid.org/0000-0002-2301-7089>

Alfredo P. Santos Jr  <https://orcid.org/0000-0002-2829-718X>

Isabelly G. Martins  <https://orcid.org/0009-0002-4420-5713>

Roberta Graboski  <https://orcid.org/0000-0002-9123-4819>

Wilian Vaz-Silva  <https://orcid.org/0000-0001-6235-5331>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Almeida JPFA, Freitas MADE, Silva MB, Valverde MCC, Rodrigues MT, Pires AM, Mott T (2018) A new four-pored *Amphisbaena* (Squamata: Amphisbaenidae) from northeastern Brazil. *Zootaxa* 4514(4): 553–562. <https://doi.org/10.11646/zootaxa.4514.4.8>
- Angiolani-Larrea F, de Fraga R, Mortati A, Ribeiro S (2021) Sex-dependent divergence in asymmetrical reproductive tract morphology between two closely-related amphisbaenian species (Squamata: Amphisbaenidae). *Amphibia-Reptilia* 42(3): 317–330. <https://doi.org/10.1163/15685381-bja10056>
- Costa HC (2020) New record and updated distribution map of the rare *Amphisbaena spurrelli* (Amphisbaenia: Amphisbaenidae). *Phyllomedusa* 19(1): 259–266. <https://doi.org/10.11606/issn.2316-9079.v19i2p259-266>
- Costa HC, Resende FC, Teixeira Jr M, Dal Vechio F, Clemente CA (2015) A new *Amphisbaena* (Squamata: Amphisbaenidae) from southern Espinhaço Range, southeastern Brazil. *Anais da Academia Brasileira de Ciências* 87(2): 891–901. <https://doi.org/10.1590/0001-3765201520140088>
- Costa HC, Welton LJ, Hallermann J (2018) An updated diagnosis of the rare *Amphisbaena slateri* Boulenger, 1907, based on additional specimens (Squamata, Amphisbaenia, Amphisbaenidae). *Evolutionary Systematics* 2(2): 125–135. <https://doi.org/10.3897/evolsyst.2.28059>
- Costa HC, Velasquez S, Zaher H, Garcia PCA (2019) Updated diagnosis of *Amphisbaena metallurga* and *A. sanctaeritae* and first record of *A. hiata* in Brazil (Squamata: Amphisbaenidae). *South American Journal of Herpetology* 14(3): 233–241. <https://doi.org/10.2994/SAJH-D-17-00105.1>

- Cotrim CGC (2015) “As bellas amethystas”: garimpos e relações comerciais em Brejinho das Ametistas (Alto Sertão da Bahia, 1870–1930). *Epígrafe* 2(2): 25–46. <https://doi.org/10.11606/issn.2318-8855.v2i2p25-46>
- Dal Vechio F, Teixeira Jr M, Mott T, Rodrigues MT (2018) Rediscovery of the poorly known *Amphisbaena bahiana* Vanzolini, 1964 (Squamata, Amphisbaenidae), with data on its phylogenetic placement, external morphology and natural history. *South American Journal of Herpetology* 13(3): 238–248. <https://doi.org/10.2994/SAJH-D-16-00053.1>
- Dias AMS, Fonseca A, Paglia AP (2019) Technical quality of fauna monitoring programs in the environmental impact assessments of large mining projects in southeastern Brazil. *Science of the Total Environment* 650(1): 216–223. <https://doi.org/10.1016/j.scitotenv.2018.08.425>
- Gans C (1961) Notes on amphisbaenids (Amphisbaenia: Reptilia). 2. *Amphisbaena occidentalis* Cope from the coastal plain of northern Peru. *Postilla* (56): 1–17.
- Gans C (1962a) Notes on amphisbaenids (Amphisbaenia: Reptilia). 5. A redefinition and a bibliography of *Amphisbaena alba* Linné. *American Museum Novitates* 2105: 1–31.
- Gans C (1962b) Notes on amphisbaenids (Amphisbaenia: Reptilia). 6. Redescription and range extension of *Amphisbaena spurrelli* Boulenger. *Breviora* 171: 1–11.
- Gans C (1963a) Notes on amphisbaenids (Amphisbaenia: Reptilia). 10. Redescription and redefinition of *Amphisbaena pericensis* Noble from the mountains of northwestern Peru. *Breviora* 189: 1–15.
- Gans C (1963b) Notes on amphisbaenids (Amphisbaenia: Reptilia). 9. Redescription of *Amphisbaena ridleyi* Boulenger. *Copeia* 1963(1): 102–107. <https://doi.org/10.2307/1441276>
- Gans C (1963c) Notes on amphisbaenids (Amphisbaenia, Reptilia). 8. A redescription of *Amphisbaena stejnegeri* and the description of a new species of *Amphisbaena* from British Guiana. *American Museum Novitates* 2128: 1–18.
- Gans C (1964a) Notes on amphisbaenids (Amphisbaenia: Reptilia). 11. On *Amphisbaena schmidtii*, a third species of the genus from Puerto Rico (Amphisbaenia: Reptilia). *Breviora* 198: 1–11.
- Gans C (1964b) On the South American species of *Amphisbaena* with a vertically keeled tail (Amphisbaenia: Reptilia). Notes on amphisbaenids. 15. *Senckenbergiana biologia* (Mertens Festschrift) 45(3/5): 387–416.
- Gans C (1965a) Notes on amphisbaenids. 17. Redescription and discussion of *Amphisbaena angustifrons* Cope and *A. camura* Cope, large amphisbaenids of southern South America (Amphisbaenia: Reptilia). *American Museum Novitates* (2225): 1–32.
- Gans C (1965b) Notes on amphisbaenids (Amphisbaenia: Reptilia). 16. On *Amphisbaena heathi* Schmidt and *A. carvalhoi* new species, small forms from the northeast of Brazil (Amphisbaenia: Reptilia). *Proceedings of the California Academy of Science* 31(23): 613–630.
- Gans C (1966) Studies on amphisbaenids (Amphisbaenia: Reptilia) 3. The small species from southern South America commonly identified as *Amphisbaena darwini*. *Bulletin of the American Museum of Natural History* 134(3): 185–260.
- Gans C (1967) Redescription of *Amphisbaena slateri* Boulenger, with comments on its range extension into Bolivia. *Herpetologica* 23(3): 223–227.
- Gans C, Alexander AA (1962) Studies on the amphisbaenids (Amphisbaenia; Reptilia). 2. On the amphisbaenids of the Antilles. *Bulletin of the Museum of Comparative Zoology* 128(3): 65–158.
- Gans C, Amdur M (1966) Redescription of *Amphisbaena vermicularis* Wagler, with comments on its range and synonymy (Amphisbaenia: Reptilia). Notes on amphisbae-

- nids 20. *Proceedings of the California Academy of Science* 33(5): 69–90. <https://doi.org/10.2307/1441079>
- Gans C, Diefenbach COC (1972) Description and geographical variation in the South American *Amphisbaena angustifrons*; the southernmost amphisbaenian in the world (Reptilia: Amphisbaenia). *American Museum Novitates* 2494: 1–20.
- Gans C, Mather S (1977) *Amphisbaena medemi*, an interesting new species from Colombia (Amphisbaenia, Reptilia), with a key to the amphisbaenians of the Americas. *Fieldiana (Zoology)* 72(2): 21–46. <https://doi.org/10.5962/bhl.title.5131>
- Gans C, Rhodes C (1964) Notes on amphisbaenids (Amphisbaenia: Reptilia). 13. A systematic review of *Anops* Bell, 1833. *American Museum Novitates* 2186: 1–25.
- Gil-Jiménez E, Lucas M, Ferrer M (2021) Metalliferous mining pollution and its impact on terrestrial and semi-terrestrial vertebrates: a review. In: Voogt P (Ed.) *Reviews of environmental contamination and toxicology* 256: 1–69. Springer. https://doi.org/10.1007/398_2021_65
- González-Sponga MA, Gans C (1971) *Amphisbaena gracilis* Strauch. Rediscovered (Amphisbaenia: Reptilia). *Copeia* 1971(4): 589–595. <https://doi.org/10.2307/1442627>
- Graboski R., Grazziotin FG, Mott T, Rodrigues MT (2022) The phylogenetic position of Ridley’s worm lizard reveals the complex biogeographic history of New World insular amphisbaenids. *Molecular Phylogenetics and Evolution* 173: 107518. <https://doi.org/10.1016/j.ympev.2022.107518>
- Graboski R, Arredondo JC, Grazziotin FG, Guerra-Fuentes RA, Da Silva AAA, Prudente ALC, Pinto RR, Rodrigues MT, Bonatto SL, Zaher H (2023) Revealing the cryptic diversity of the widespread and poorly known South American blind snake genus *Amerotyphlops* (Typhlopidae: Scolecophidia) through integrative taxonomy. *Zoological Journal of the Linnean Society* 197: 719–751. <https://doi.org/10.1093/zoolinnean/zlac059>
- Grillitsch B, Schiesari L (2010) The ecotoxicology of metals in reptiles In: Sparling DW, Linder G, Bisho CA, Krest S (Eds) *Ecotoxicology of amphibians and Reptiles*, 2nd edn.: 337–448. CRC Press, New York, 944 pp. <https://doi.org/10.1201/EBK1420064162-c12>
- Guedes TB, Entiauspe-Neto OM, Costa HC (2023) Lista de répteis do Brasil: atualização de 2022. *Herpetologia Brasileira* 12(1): 56–161. <https://doi.org/10.5281/zenodo.7829013>
- Hoogmoed MS (1989) A new species of *Amphisbaena* (Amphisbaenia: Amphisbaenidae) from Suriname. Notes on the herpetofauna of Surinam XI in honour of Dr. Pieter Wagenaar Hummelinck. *Foundation for Scientific Research in Surinam and the Netherlands Antilles (Amsterdam)* 123: 65–73.
- Hoogmoed MS, Ávila-Pires TCS (1991) A new species of small *Amphisbaena* (Reptilia: Amphisbaenia: Amphisbaenidae) from western Amazonian Brazil. *Boletim do Museu Paraense Emilio Goeldi Série Zoologia* 7(1): 77–94.
- IBGE (2023) Banco de Informações Ambientais. <https://www.ibge.gov.br/geociencias/informacoes-ambientais/geologia/23382-banco-de-informacoes-ambientais.html> [accessed 18 November 2023]
- Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kearney M, Stuart BL (2004) Repeated evolution of limblessness and digging heads in worm lizards revealed by DNA from old bones. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 271(1549): 1677–1683. <https://doi.org/10.1098/rspb.2004.2771>
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2016) PartitionFinder 2: New methods for selecting partitioned models of evolution for molecular and

- morphological phylogenetic analyses. *Molecular Biology and Evolution* 34(3): 772–773. <https://doi.org/10.1093/molbev/msw260>
- Longrich NR, Vinther J, Pyron RA, Pisani D, Gauthier JA (2015) Biogeography of worm lizards (Amphisbaenia) driven by end-Cretaceous mass extinction. *Proceedings of the Royal Society B: Biological Sciences* 282(1806): 21396–21401. <https://doi.org/10.1098/rspb.2014.3034>
- Montero R (1996) *Amphisbaena bolivica* Mertens 1929, nueva combinación (Squamata: Amphisbaenia). *Cuadernos de Herpetología* 9(2): 75–84.
- Montero R, Sáfadez IF, González Álvarez L (1997) On a new species of *Amphisbaena* (Reptilia: Amphisbaenidae) from Bolivia. *Journal of Herpetology* 31(2): 218–220. <https://doi.org/10.2307/1565389>
- Mott T, Vieites DR (2009) Molecular phylogenetics reveals extreme morphological homoplasy in Brazilian worm lizards challenging current taxonomy. *Molecular Phylogenetics and Evolution* 51(2): 190–200. <https://doi.org/10.1016/j.ympev.2009.01.014>
- Mott T, Rodrigues MT, Freitas MA, Silva TFS (2008) New species of *Amphisbaena* with a non-autotomic and dorsally tuberculate blunt tail from State of Bahia, Brazil (Squamata, Amphisbaenidae). *Journal of Herpetology* 42(1): 172–175. <https://doi.org/10.1670/07-074R2.1>
- Mott T, Rodrigues MT, Santos EM (2009) A new *Amphisbaena* with chevron-shaped anterior body annuli from state of Pernambuco: Brazil (Squamata: Amphisbaenidae). *Zootaxa* 2165(1): 52–58. <https://doi.org/10.11646/zootaxa.2165.1.4>
- Mott T, Carvalho-Neto CS, Carvalho-Filho KS (2011) *Amphisbaena miringoera* Vanzolini, 1971 (Squamata: Amphisbaenidae): New state record. *Check List* 7(5): 594–595. <https://doi.org/10.15560/7.5.594>
- Oliveira ECS, Vaz-Silva W, Santos-Jr AP, Graboski R, Teixeira Jr M, Dal Vechio F, Ribeiro S (2018) A new four-pored *Amphisbaena* Linnaeus, 1758 (Amphisbaenia, Amphisbaenidae) from Brazilian Amazon. *Zootaxa* 4420(4): 451–474. <https://doi.org/10.11646/zootaxa.4420.4.1>
- Perez R, Borges-Martins M (2019) Integrative taxonomy of small worm lizards from Southern South America, with description of three new species (Amphisbaenia: Amphisbaenidae). *Zoologischer Anzeiger* 283(58): 124–141. <https://doi.org/10.1016/j.jcz.2019.09.007>
- Perez R, Ribeiro S, Borges-Martins M (2012) Reappraisal of the taxonomic status of *Amphisbaena prunicolor* (Cope 1885) and *Amphisbaena albocingulata* Boettger 1885 (Amphisbaenia: Amphisbaenidae). *Zootaxa* 3550(1): 1–25. <https://doi.org/10.11646/zootaxa.3550.1.1>
- Ribeiro S, Eleutério JMP (2023) Anfisbênios dos Brasil. In: Silva-Soares T, Monjardim M, Figueiredo RG (Eds) *Herpetologia para todos*. Instituto Biodiversidade Neotropical, Espírito Santo, 93–105.
- Ribeiro S, Vaz-Silva W, Santos-Jr AP (2008) New pored *Leposternon* (Squamata, Amphisbaenia) from Brazilian Cerrado. *Zootaxa* 1930(1): 18–38. <https://doi.org/10.11646/zootaxa.1930.1.2>
- Ribeiro S, Castro-Mello C, Nogueira C. (2009) New species of *Anops* Bell, 1893, (Squamata, Amphisbaenia). *Journal of Herpetology* 43(1): 21–28. <https://doi.org/10.1670/07-299R1.1>
- Ribeiro S, Nogueira C, Cintra CE, Silva Jr NJ, Zaher H (2011) Description of a new pored *Leposternon* (Squamata, Amphisbaenidae) from the Brazilian Cerrado. *South American Journal of Herpetology* 6(3): 177–188. <https://doi.org/10.2994/057.006.0303>

- Ribeiro S, Santos AP, Zaher H (2015) A new species of *Leposternon* Wagler, 1824 (Squamata, Amphisbaenia) from northeastern Argentina. *Zootaxa* 4034(2): 309–324. <https://doi.org/10.11646/zootaxa.4034.2.4>
- Ribeiro S, Gomes JO, Silva HLR, Cintra CED, Silva Jr N (2016) A new two-pored species of *Amphisbaena* (Squamata, Amphisbaenidae) from the Brazilian Cerrado, with a key to the two-pored species of *Amphisbaena*. *Zootaxa* 4147(2): 124–142. <https://doi.org/10.11646/zootaxa.4147.2.2>
- Ribeiro S, Silveira A, Santos-Jr AP (2018a) A new species of *Leposternon* (Squamata: Amphisbaenidae) from Brazilian Cerrado with a key to pored species. *Journal of Herpetology* 52(1): 50–58. <https://doi.org/10.1670/16-125>
- Ribeiro LB, Gomides SC, Costa HC (2018b) A new species of *Amphisbaena* from northeastern Brazil (Squamata: Amphisbaenidae). *Journal of Herpetology* 52(2): 234–241. <http://doi/full/https://doi.org/10.1670/17-028>
- Ribeiro S, Sá V, Santos-Jr AP, Graboski R, Zaher H, Guedes AG, Andrade SP, Vaz-Silva W (2019) A new species of the *Amphisbaena* (Squamata, Amphisbaenidae) from the Brazilian Cerrado with a key for the two-pored species. *Zootaxa* 4550(3): 301–320. <https://doi.org/10.11646/zootaxa.4550.3.1>
- Ribeiro LB, Gomides SC, Costa HC (2020) A new worm lizard species (Squamata: Amphisbaenidae: *Amphisbaena*) with non-autotomic tail, from Northeastern Brazil. *Journal of Herpetology* 54(1): 9–18. <https://doi.org/10.1670/19-043>
- Ribeiro-Júnior MA, Ribeiro S, Cintra CED, Gomes JO (2022) *Amphisbaena ibijara* Rodrigues, Andrade & Lima, 2003 is a junior synonym of *Amphisbaena frontalis* Vanzolini, 1991 (Squamata, Amphisbaenia). *Journal of Herpetology* 56(2): 234–240. <https://doi.org/10.1670/21-039>
- Roberto IJ, Brito LBM, Ávila RW (2014) A new six-pored *Amphisbaena* (Squamata: Amphisbaenidae) from the coastal zone of northeast Brazil. *Zootaxa* 3753(2): 167–176. <https://doi.org/10.11646/zootaxa.3753.2.6>
- Rocha CFD, Barros-Filho JD, Sluys MV (2023) A new *Amphisbaena* species from a restinga habitat of Southeastern Brazil (Reptilia, Squamata, Amphisbaenidae). *International Journal of Zoology and Animal Biology* 6(4): 1–8. <https://doi.org/10.23880/izab-16000501>
- Rodrigues MT, Andrade GV, Lima JD (2003) A new species of *Amphisbaena* (Squamata, Amphisbaenidae) from state of Maranhão, Brazil. *Phyllomedusa* 2(1): 21–26. <https://doi.org/10.11606/issn.2316-9079.v2i1p21-26>
- Sindaco R, Panagiotis K, Roberto S, Petros L (2014) Taxonomic reassessment of *Blanus strauchi* (Bedriaga, 1884) (Squamata: Amphisbaenia: Blanidae), with the description of a new species from south-east Anatolia (Turkey). *Zootaxa* 3795(3): 311–326. <https://doi.org/10.11646/zootaxa.3795.3.6>
- Spix JB, Martius KFP (1938) *Através da Bahia: Excerptos da obra Reise in Brasilien*. Traduzidos a português pelos Drs. Pirajá da Silva e Paulo Wolf. Companhia Editora Nacional, São Paulo, 342 pp.
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Strüssmann C, Carvalho MA (2001) Two new species of *Cercolophia* Vanzolini, 1992 from the state of Mato Grosso, western Brazil (Reptilia, Amphisbaenia, Amphisbaenidae). *Bollettino del Museo Regionale di Scienze Naturali di Torino* 18(2): 487–505.

- Swofford DL (2003) PAUP*: Phylogenetic Analysis Using Parsimony (*and Other Methods). Sunderland, Massachusetts: Sinauer Associates. <https://paup.phylosolutions.com> [accessed 08 December 2023]
- Teixeira Jr M, Dal Vechio F, Neto AM, Rodrigues MT (2014) A new two-pored *Amphisbaena* Linnaeus, 1758, from Western Amazonia, Brazil (Amphisbaenia: Reptilia). *South American Journal of Herpetology* 9(1): 62–74. <https://doi.org/10.2994/SA-JH-D-14-00004.1>
- Teixeira Jr M, Dal Vechio F, Rodrigues MT (2016) Diagnostic clarification, new morphological data and phylogenetic placement of *Amphisbaena arenaria* Vanzolini, 1991 (Amphisbaenia, Amphisbaenidae). *Zootaxa* 4205(3): 293–296. <https://doi.org/10.11646/zootaxa.4205.3.9>
- Teixeira Junior M, Dal Vechio F, Recoder R, Cassimiro J, Sena MA, Rodrigues MT (2019) Two new highland species of *Amphisbaena* Linnaeus, 1758 (Amphisbaenia, Amphisbaenidae) from Bahia State, Brazil. *South American Journal of Herpetology* 14(3): 213–232. <https://doi.org/10.2994/SAJH-D-17-00097.1>
- Thomas R (1965) Two new subspecies of *Amphisbaena* (Amphisbaenia, Reptilia) from the Barahona Peninsula of Hispaniola. *Breviora* 215: 1–14.
- Thomas R (1966) Additional notes on the amphisbaenids of Greater Puerto Rico. *Breviora* 249: 1–23.
- Thomas R, Hedges SB (1998) A new amphisbaenian from Cuba. *Journal of Herpetology* 32(1): 92–96. <https://doi.org/10.2307/1565485>
- Thomas R, Hedges SB (2006) Two new species of *Amphisbaena* (Reptilia: Squamata: Amphisbaenidae) from the Tiburon Peninsula of Haiti. *Caribbean Journal of Science* 42(2): 208–219.
- Torres-Ramírez JJ, Angarita-Sierra T, Vargas-Ramírez M (2021) A new species of *Amphisbaena* (Squamata: Amphisbaenidae) from the Orinoquian region of Colombia. *Vertebrate Zoology* 71(274): 55–74. <https://doi.org/10.3897/vz.71.e59461>
- Vanzolini PE (1950) Contribuições ao conhecimento dos lagartos brasileiros da família Amphisbaenidae Gray, 1825. I. Sobre uma nova subespécie insular de *Amphisbaena darwini* D. and B., 1839. *Papéis Avulsos de Zoologia* 9(6): 69–78. <https://doi.org/10.11606/0031-1049.1950.9p69-78>
- Vanzolini PE (1991) Two further new species of *Amphisbaena* from the semi-arid north-east of Brasil (Reptilia, Amphisbaenia). *Papéis Avulsos de Zoologia* 37(23): 347–361. <https://doi.org/10.11606/0031-1049.1990.37.p347-361>
- Vanzolini PE (1992) *Cercolophia*, a new genus for the species of *Amphisbaena* with a terminal vertical keel on the tail. *Papéis Avulsos de Zoologia* 37(27): 401–412. <https://doi.org/10.11606/0031-1049.1990.37.p401-412>
- Vanzolini PE (1995) A new species of *Amphisbaena* from the state of Mato Grosso, Brazil (Squamata, Amphisbaenidae). *Papéis Avulsos de Zoologia* 3(10): 217–221. <https://doi.org/10.11606/0031-1049.1994.39.p217-221>
- Vanzolini PE (1996) On slender species of *Amphisbaena*, with the description of a new one from northeastern Brasil (Squamata, Amphisbaenia). *Papéis Avulsos de Zoologia* 39(16): 293–305. <https://doi.org/10.11606/0031-1049.1994.39.p293-305>
- Vanzolini PE (2002) An aid to the identification of the South American species of *Amphisbaena* (Squamata, Amphisbaenidae). *Papéis Avulsos de Zoologia* 42(15): 351–362. <https://doi.org/10.1590/S0031-10492002001500001>
- Vaz-Silva W (2009) Herpetofauna im Zentralbrasilianischen Cerrado. *Terraria (Munster)* 17: 75–78.

Supplementary material 1

Supplementary information

Authors: Síria Ribeiro, Alfredo P. Santos Jr, Isabelly G. Martins, Elaine C. S. Oliveira, Roberta Graboski, Thiago Barbosa Da Silveira, Matheus H. M. Benício, Wilian Vaz-Silva

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1213.122265.suppl1>